

# Marin County Wildland Fires

Examining Fuel Load and Land Cover Change to Inform Fire  
Prevention and Suppression Decisions in Marin County, CA

Suhani Dalal <sup>xx</sup> Katera Lee <sup>xx</sup> Chandler Ross <sup>xx</sup> Gabriel Rosenstein


# Marin County, CA

## Marin County

Located between the Pacific Ocean and the San Francisco Bay, directly north of San Francisco

## Study Period

May – November; 2018 – 2022



Basemap: 2022 PlanetScope Imagery




# Project Partners



**Marin  
County  
Fire**

**Department**



**FIRE  
Foundry**





# Community Concerns

- People in surrounding areas are at **higher risk**
- Proximity to infrastructure damage
- **Destruction** of natural areas



Source: Tony Webster, Wikimedia Commons





# Objectives



## Fuel Load Maps

Use various vegetation indices to determine the load and density of fuel for wildfires



# Objectives



## Fuel Load Maps



## Land Cover Maps

Identify the aspect, slope, elevation, and land cover types across study period



# Objectives



## Fuel Load Maps



## Land Cover Maps



## Vegetation Moisture Maps

Utilize high resolution ECOSTRESS data to examine soil and vegetation moisture parameters





# Objectives



## Fuel Load Maps



## Land Cover Maps




## Vegetation Moisture Maps



## Fire Suppression Model

Input all factors associated with wildfire risk severity into a complete model in ArcGIS Pro that can guide fire suppression decision making

# Earth Observation Platforms and Sensors




*International Space Station  
(ISS)- ECOSTRESS*



*Landsat 7 ETM+ and  
Landsat 8 OLI & TIRS*



*Sentinel 2-A MSI*









*PlanetScope DOVE*



*LiDAR*


# Data Processing: Topography





# Data Processing: Moisture

## ECOSTRESS:



# Data Processing: Landcover Fuels


Gather  
Sentinel-2A  
images



Run Dynamic  
World code in  
GEE




Process annual  
composites in  
ArcGIS





Wet season NDVI  
(mid-November  
through mid-May)



Dry season NDVI  
(mid-May  
through mid-




=




NDVI differential (wet season - dry season)


# Data Processing: Forest Fuels




Canopy Bulk  
Density




Canopy Base  
Height



Canopy Height




Canopy Cover





Ladder Fuel  
Density

Ladder Fuel  
Density




Pre-fire ladder  
fuels



Fire perimeter  
burn severity



Ladder Fuel  
Density




Post-fire ladder  
fuels




# Fuel

S





## Ladder Fuel Density



## Canopy Cover



## Landcover




## Canopy Base Height


## Canopy Bulk Density

## NDVI Differential

# Data Analysis




# Model 1: FlamMap






# Model 2: Suitability



# Model 3: Machine Learning






# Methodology: Fire Lines




Image Credit: A. J. T. Johnsingh, WWF-India and NCF






# Results: Severity Models

## FlamMap Severity




## Suitability Severity






# Results: Severity Models


## FlamMap Severity



## Suitability Severity





## Machine Learning





# Results: Fireline Model



1 Mile




- Roads
- Trails
- Water

## Fireline Potential

- Easy
- Medium
- Difficult

# Errors and Uncertainties



Data Limitations



Model Uncertainties





# Conclusions

- A fusion of satellite and ground LiDAR data, focused on moisture, fuels, and topography can be used to quantify fire severity in Marin's unique environment of microclimates
- Model comparisons can be used to understand the various algorithms for computing fire severity in relation to a region
- A feasible approach is combining the outputs of the fire severity models with fire line & slope data for the fire line model



Image Credit: Fire Foundry





# Future Work



**Locate** areas for fire lines beyond Marin County



**Validate** input parameters



**Incorporate** additional parameters into the fire model



**Design** a workshop for FIRE Foundry



# Acknowledgements

## Partners

- **FIRE Foundry**
  - Josh Dimon
- **Marin County Fire Department**
  - Graham Groneman

## NASA DEVELOP

- Lisa Tanh (NASA DEVELOP Ames Research Center Fellow)

## Science Advisors

- Dr. Jake Dialesandro (Santa Clara University)
- Dr. Juan Torres-Pérez (NASA Ames Research Center)
- Britnay Beaudry (NASA Ames Research Center)

Includes copyrighted material of Planet Labs PBC. All rights reserved.

This work utilized data made available through the NASA Commercial Smallsat Data Acquisition (CSDA) program.

This material is based upon work supported by NASA through contract NNL16AA05C. Any mention of a commercial product, service, or activity in this material does not constitute NASA endorsement. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Aeronautics and Space Administration and partner organizations.